

*Please store this instruction leaflet in a safe place for later reference.*

*Be sure to read the instructions carefully before use.*

## HIYA ROKUSHINGAN

樋屋六神丸

<CARDIOTONIC>

“HIYA ROKUSHINGAN”, containing highly-selected crude drug, show the excellent efficacy on the palpitation and panting.

### CAUTIONS FOR USE

*[To be avoided!]*

(Failure to observe the following may result in the present symptoms worsening or adverse reactions/accidents occurring.)

This medication should not be used in combination with the following medications:

Any other cardiotonics

*[Consult a physician for the following]*

1. The following persons should consult a physician, pharmacist, or registered distributor of this medication before using this medication.
  - (1)A person undergoing medical treatment from a physician
  - (2)A pregnant woman or a woman suspected of being pregnant
2. The following symptoms may be adverse reactions to this medication. If any of these symptoms occur, discontinue use immediately and consult a physician, pharmacist, or registered distributor of this medication. Take the instruction leaflet with you.

Related area	Symptoms
Digestive organs	Nausea, vomiting

3. Discontinue use when the cold symptoms do not improve after having used this medication for 5 or 6 days, and consult a physician, pharmacist, or registered distributor of this medication. Take the instruction leaflet with you.

### INDICATIONS

Palpitation, panting, restorative

**COMPOSITION (in 6 pills)**

Components	Content
Toad cake	4.5 mg
Musk	1.0 mg
Oriental bezoar	3.0 mg
Ginseng	4.0 mg
Animal bile	3.0 mg
<i>α</i> -Borneol	2.0 mg

Additives: Rice Powder, Kanbaiko, Lactose, Honey, Paraben, Gold Leaf

**DOSAGE AND ADMINISTRATION**

Age	One dosage	Daily dosage
15 years ~	2 pills	3 times a day before or between each meal.
Under 15 years	Do not use	

*[Cautions]*

- (1) Use this medication without chewing.
- (2) Please follow the recommended dosage and administration.

**CAUTIONS ON HANDLING AND STORAGE**

- (1) Store in a sealed container in a cool, dry place free from direct sunlight.
- (2) Keep all medication out of the reach of children.
- (3) Do not transfer this medicine to a different container. (It may be misused or the quality may deteriorate.)