

*Please store this instruction leaflet in a safe place for later reference.*

*Be sure to read the instructions carefully before use.*

## HIYA KODOMO SOGO KAZEGUSURI

ヒヤこども総合かぜ薬

<COLD MEDICATION>

“**HIYA KODOMO SOGO KAZEGUSURI**” is cold medication for the relief of symptoms associated with the common cold. Approved for use by children between 1 to 14 years of age.

### <PRODUCT FEATURES>

1. Containing kakkonto extract (Kampo-medicine) which is frequently used to treat incipient cold
2. Easy-to-take granules with chocolate flavor suitable for small children

### CAUTIONS FOR USE

(Though HIYA KODOMO SOGO KAZEGUSURI is cold medication for children, common cautions as cold medication are described here.)

*[To be avoided!]*

(Failure to observe the following may result in the cold symptoms worsening or adverse reactions/accidents occurring.)

1. This medication should not be used by the following persons:
  - (1)Persons who experience an allergic reaction to this medication, any component of this medication
  - (2)Persons who suffer from asthma after using this medication or any other cold remedies or anti-pyretics and analgesics
2. This medication should not be used in combination with the following medications:  
Oral medicines containing other cold remedies, anti-pyretics and analgesics, sedatives, cough and expectorant remedies, or antihistamines(oral medications for nasal catarrh, motion sickness, or anti-allergic agents, etc.)
3. Do not drive or operate machinery after using this medication. (It may cause drowsiness, etc.)
4. Do not consume alcohol before or after taking this medication.
5. Do not use this medication for a prolonged period of time.

*[Consult a physician for the following]*

1. The following persons should consult a physician, pharmacist, or registered distributor of this medication before using this medication.

(1) A person undergoing medical treatment from a physician or dentist

(2) A pregnant woman or a woman suspected of being pregnant

(3) An elderly person

(4) A person with a medical history of being allergic to medicines and other substances

(5) A person with the following symptoms:

High fever, difficulty in urination

(6) A person diagnosed as having the following:

Thyroid disease, diabetes mellitus, cardiac disease, hypertension, hepatic disease, renal disease, gastric/duodenal ulcer, glaucoma

2. The following symptoms may be adverse reactions to this medication. If any of these symptoms occur, discontinue use immediately and consult a physician, pharmacist, or registered distributor of this medication. Take the instruction leaflet with you.

Related area	Symptoms
Skin	Rash, redness, itching
Digestive organs	Nausea, vomiting, loss of appetite
Central nervous system	Dizziness
Urinary system	Difficulty in urination
Other	Significant decrease in temperature

In very rare cases the following serious symptoms may occur. In this case, consult a physician immediately.

Symptom's name	Symptoms
Shock(anaphylaxis)	Occurrence of itching of skin, hives, hoarseness, sneezing, itching of throat, breathing difficulties, palpitations, or clouding of consciousness, etc. soon after use
Muco-cutaneo-ocular syndrome(Stevens-Johnson syndrome), toxic epidermal necrolysis, acute generalized exanthematous pustulosis	Continuation or sudden worsening of the following: high fever, redness of the eye, eye discharge, erosion of the lip, throat pain, rash or redness over an extensive area of skin, red circular skin bumps(pustules), general listlessness, loss of appetite, etc.

Hepatic function failure	Fever, itching, rash, jaundice(the skin and whites of the eyes turn yellow), brown urine, general listlessness, loss of appetite, etc.
Interstitial pneumonia	Sudden occurrence or continuation of the following: shortness of breath or choking sensation after going up stairs or following exertion, dry cough, fever, etc.
Asthma	Wheezing and labored breathing
Aplastic anemia	Bruising, nosebleed, gingival bleeding, fever, pale skin and mucus, fatigue, palpitation, shortness of breath, feel ill and dizzy, and appearance of blood in urine, etc.
Agranulocytosis	Sudden high fever, chill, throat pain, etc.
Renal impairment	Decreased urinary output, fever, rash, general swelling, general listlessness, joint pain, diarrhea, etc.

3. The following symptoms may appear with use of this medication. If these symptoms continue or worsen, discontinue use and consult a physician, pharmacist, or registered distributor of this medication. Take the instruction leaflet with you.

Dry mouth, drowsiness

4. Discontinue use when the cold symptoms do not improve after having used this medication 5 or 6 times, and consult a physician, pharmacist, or registered distributor of this medication. Take the instruction leaflet with you.

## INDICATIONS

Relief of cold symptoms (a runny nose, nasal congestion, sneezing, coughs, sputum, sore throat, chills, fever, headaches, joint pain, and muscular pain)

## COMPOSITION (in 6 sachets/4.8g)

Components	Content
Acetaminophen	300 mg
Tipepidine hibenzate	37.5 mg
Chlorpheniramine maleate	3.75 mg
Caffeine and sodium benzoate	100 mg
Kakkonto extract	1000 mg (Extracted from 1.333 g of Pueraria root, 0.667 g of Ephedra herb, 0.667 g of Jujube, 0.5 g of Cinnamon bark, 0.5 g of Peony root, 0.333 g of Glycyrrhiza, 0.167 g of Ginger)

Additives: Sucrose, Lactose, Magnesium stearate, Magnesium aluminometasilicate, flavor, Ethyl vanillin, Vanillin, and Propylene glycol

#### **DOSAGE AND DIRECTION**

Age	One dosage	Daily dosage
7 up to 15 years	2 sachets	3 times daily within 30 minutes after eating.
3 up to 7 years	1 sachet	
1 up to 3 years	3/4 sachet	
Under 1 year	Do not use	

#### *[Cautions]*

- (1) This medication should be given to children only under adult supervision.
- (2) For children aged 2 years or younger, make it a priority to consult a physician. Use this medication only in case it's really necessary.
- (3) Please follow the recommended dosage and directions.

#### **CAUTIONS ON HANDLING AND STORAGE**

- (1) Store in a sealed container in a cool, dry place free from direct sunlight.
- (2) Keep all medication out of the reach of children.
- (3) Do not transfer this medicine to a different container. (It may be misused or the quality may deteriorate.)
- (4) Do not exceed the period of use.