

*Please store this instruction leaflet in a safe place for later reference.*

*Be sure to read the instructions carefully before use.*

## HIYA KODOMO SEKI SYRUP N

ヒヤこどもせきシロップ N

<COUGH MEDICATION>

Designated Class 2 OTC drug

“HIYA KODOMO SEKI SYRUP N” is cough medication for the relief of cough and sputum of children caused by various reasons. Approved for use by children aged 3 months up to 15 years.

<PRODUCT FEATURES>

1. Sugar-free (containing xylitol)
2. Caffeine-free
3. Easy-to-drink syrup with muskmelon flavor suitable for small children
4. Colorant-free
5. Child resistant cap

### CAUTIONS FOR USE

{Though HIYA KODOMO SEKI SYRUP N is cough medication for children, common cautions as cough and expectorant remedies (cough medication) are described here.}

*[To be avoided!]*

(Failure to observe the following may result in the symptoms worsening or adverse reactions/accidents occurring.)

1. This medication should not be used in combination with the following medications:  
Oral medicines containing other cough and expectorant remedies, cold remedies, sedatives, or antihistamines (oral medications for nasal catarrh, motion sickness, or anti-allergic agents, etc.)
2. Do not drive or operate machinery after using this medication. (It may cause drowsiness, etc.)
3. Nursing women should either not take this medication or stop nursing while taking while taking the medication.
4. Do not use this medication for a prolonged period of time, nor overdose.

*[Consult a physician for the following]*

1. The following persons should consult a physician, pharmacist, or registered distributor of this medication before using this medication.

(1) A person undergoing medical treatment from a physician

(2) A pregnant woman or a woman suspected of being pregnant

(3) An elderly person

(4) A person with a medical history of being allergic to medicines and other substances

(5) A person with the following symptoms:

High fever, difficulty in urination

(6) A person diagnosed as having the following:

Cardiac disease, hypertension, diabetes mellitus, glaucoma, thyroid disease

2. The following symptoms may be adverse reactions to this medication. If any of these symptoms occur, discontinue use immediately and consult a physician, pharmacist, or registered distributor of this medication. Take the instruction leaflet with you.

Related area	Symptoms
Skin	Rash, redness, itching
Central nervous system	Dizziness
Digestive organs	Nausea, vomiting, loss of appetite
Urinary system	Difficulty in urination

In very rare cases the following serious symptoms may occur. In this case, consult a physician immediately.

Symptom's name	Symptoms
Aplastic anemia	Bruising, nosebleed, gingival bleeding, fever, pale skin and mucus, fatigue, palpitation, shortness of breath, feel ill and dizzy, and appearance of blood in urine, etc.
Agranulocytosis	Sudden high fever, chill, throat pain, etc.

3. The following symptoms may appear with use of this medication. If these symptoms continue or worsen, discontinue use and consult a physician, pharmacist, or registered distributor of this medication. Take the instruction leaflet with you.

Constipation, dry mouth, drowsiness

4. Discontinue use when the cold symptoms do not improve after having used this medication 5 or 6 times, and consult a physician, pharmacist, or registered distributor of this medication. Take the instruction leaflet with you.

## INDICATIONS

Cough and sputum

**COMPOSITION (in 36 mL)**

Components	Content
Tipevidin citrate	36 mg
<i>d</i> -Methylephedrine hydrochloride	45 mg
Potassium guaiacolsulfonate	162 mg
Chlorpheniramine maleate	7.2 mg

Additives: Xylitol, D-sorbitol, Sodium chloride, Citric acid, Sodium citrate, Saccharin sodium, Sodium benzoate, Paraben, Propylene glycol, and flavor

**DOSAGE AND DIRECTION**

Age	One dosage	Daily dosage
11 up to 15 years	6 mL	3 times daily after eating, and before bedtime. Maximum 6 times daily, with an interval of 4 hours.
8 up to 11 years	5 mL	
5 up to 8 years	3 mL	
3 up to 5 years	2.5 mL	
1 up to 3 years	2 mL	
3 months up to 1 year	1 mL	
Under 3 months	Do not use	

*[Cautions]*

- (1) Please follow the recommended dosage and directions.
- (2) This medication should be given to children only under adult supervision.
- (3) For children aged 2 years or younger, make it a priority to consult a physician. Use this medication only in case it's really necessary.

**CAUTIONS ON HANDLING AND STORAGE**

- (1) Store in a sealed container in a cool, dry place free from direct sunlight.
- (2) Keep all medication out of the reach of children.
- (3) Do not transfer this medicine to a different container. (It may be misused or the quality may deteriorate.)
- (4) Do not exceed the period of use.
- (5) Do not store for long periods after opening the container. (To ensure the quality of the medication)